To be sold by retail on the prescription of a Registered Medical Practitioner only.

MSN

PRESCRIBING INFORMATION

1. GENERIC NAME

Voriconazole 50 mg and 200 mg film-coated tablets
Voriconazole 200 mg powder for solution for infusion

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Voriconazole tablets
Each Film Coated Tablet Contains
Voriconazole Tablets ——— 50 mg
Voriconazole tablets
Each Film Coated Tablet Contains
Voriconazole Tablets ——— 200 mg
Voriconazole powder for solution for infusion
Each Vial contains
Voriconazole ——— 200 mg
(As sterile freeze dried powder for reconstitution)

3. DOSAGE FORM AND STRENGTH

Voriconazole 50 mg and 200 mg film-coated tablets and Voriconazole 200 mg powder for solution for infusion.

4. CLINICAL PARTICULARS

4.1. Indications

Voriconazole is indicated for the treatment of invasive aspergillosis and serious fungal infections caused by *Scedosporium apiospermum* (asexual form of *Pseudallescheria boydii*) and *Fusarium* spp. including *Fusarium solani*, in patients intolerant of, or refractory to, other therapy.
Voriconazole is also indicated for the treatment of candidaemia in non-neutropenic patients and the following candida infections: disseminated infections in skin and infections in abdomen, kidney, bladder wall, and wounds and esophageal candidiasis.
Voriconazole is indicated for treatment of fluconazole-resistant serious invasive Candida infections (including *C. krusei*).
Voriconazole should be administered primarily to patients with proven/serious, possibly life-threatening infections.
Prophylaxis of invasive fungal infections in high risk allogeneic hematopoietic stem cell transplant (HSCT) recipients

4.2. Posology and Method of Administration

Posology
Electrolyte disturbances such as hypokalemia, hypomagnesemia and hypocalcemia should be monitored and corrected, if necessary, prior to initiation and during voriconazole therapy (see section 4.4).
It is recommended that Voriconazole is administered at a maximum rate of 3 mg/kg per hour over 1 to 3 hours.
Other brands of Voriconazole are available as 50 mg and 200 mg film-coated tablets and 40 mg/ml powder for oral suspension.

Treatment
Adults
Therapy must be initiated with the specified loading dose regimen of either intravenous or oral Voriconazole to achieve plasma concentrations on Day 1 that are close to steady state. On the basis of the high oral bioavailability (96%; see section 5.2), switching between intravenous and oral administration is appropriate when clinically indicated.
Detailed information on dosage recommendations is provided in the following table:

	Intravenous	Oral	
	Prodrug	Patients 40 kg and above*	Patients less than 40 kg*
Loading dose regimen (first 24 hours)	6 mg/kg every 12 hours	400 mg every 12 hours	200 mg every 12 hours
Maintenance dose (after first 24 hours)	4 mg/kg twice daily	200 mg twice daily	100 mg twice daily

* This also applies to patients aged 15 years and older

Duration of treatment
Treatment duration should be based on the patient's clinical and mycological response. Long term exposure to voriconazole greater than 180 days (6 months) requires careful assessment of the benefit-risk balance.

Dosage adjustment (Adults)
If patient is unable to tolerate intravenous treatment at 4 mg/kg twice daily, reduce the dose to 3 mg/kg twice daily
If patient response to treatment is inadequate, the dose may be increased to 300 mg twice daily for oral administration. For patients less than 40 kg the oral dose may be increased to 150 mg twice daily.
If patient is unable to tolerate treatment at 200 mg twice, reduce the oral dose by 50 mg steps to the 200 mg twice daily (or 100 mg twice daily for patients less than 40 kg) maintenance dose.
In case of use as prophylaxis, refer below.
Children (2 to <12 years) and young adolescents with low body weight (12 to 14 years and <50 kg)
Voriconazole should be dosed as children as these young adolescents may metabolize voriconazole more similarly to children than to adults.
The recommended dosing regimen is as follows:

	Intravenous	Oral
Loading Dose Regimen (first 24 hours)	9 mg/kg every 12 hours	Not recommended
Maintenance Dose (after first 24 hours)	8 mg/kg twice daily	9 mg/kg twice daily (a maximum dose of 350 mg twice daily)

Note: Based on a population pharmacokinetic analysis in 112 immunocompromised paediatric patients aged 2 to <12 years and 26 immunocompromised adolescents aged 12 to <17 years.
It is recommended to initiate the therapy with intravenous regimen, and oral regimen should be considered only after there is a significant clinical improvement. It should be noted that an 8 mg/kg intravenous dose in children provides exposure approximately 2-fold higher than a 9 mg/kg oral dose.
All other adolescents (12 to 14 years and ≥50 kg; 15 to 17 years regardless of body weight)
Voriconazole should be dosed as adults.
Dosage adjustment (Children [2 to <12 years] and young adolescents with low body weight [12 to 14 years and <50 kg])
If patient response to treatment is inadequate, the dose may be increased by 1 mg/kg steps. If patient is unable to tolerate treatment, reduce the intravenous dose by 1 mg/kg steps.
Use in paediatric patients aged 2 to <12 years with hepatic or renal insufficiency has not been studied (see sections 4.8 and 5.2).

Prophylaxis in Adults and Children
Prophylaxis should be initiated on the day of transplant and may be administered up to 100 days. Prophylaxis should be as short as possible depending on the risk for developing invasive aspergillosis and should be discontinued if there is anemia or immunosuppression. It may only be continued up to 180 days after transplantation in case of continuing immunosuppression or graft versus host disease (GvHD) (see section 5.1).

Dosage
The recommended dosing regimen for prophylaxis is the same as for treatment in the respective age groups. Please refer to the treatment tables above.

Duration of prophylaxis
The safety and efficacy of voriconazole use for longer than 180 days has not been adequately studied in clinical trials.
Use of voriconazole in prophylaxis for greater than 180 days (6 months) requires careful assessment of the benefit-risk balance (see sections 4.4 and 5.1).
The following instructions apply to both Treatment and Prophylaxis
Dosage adjustment
For prophylaxis use, dose adjustments are not recommended in the case of liver or renal dysfunction. However, close monitoring of these patients for treatment-related adverse events, discontinuation of voriconazole and use of alternative antifungal agents must be considered (see section 4.4 and 4.8).
Dosage adjustments in case of co-administration
Rifabutin or phenytoin may be co-administered with voriconazole if the maintenance dose of voriconazole is increased to 5 mg/kg intravenously twice daily, see sections 4.4 and 4.5.
Efavirenz may be co-administered with voriconazole if the maintenance dose of voriconazole is increased to 400 mg every 12 hours and the efavirenz dose is reduced by 50%, i.e. to 300 mg once daily. When treatment with voriconazole is stopped, the initial dosage of efavirenz should be restored (see sections 4.4 and 4.5).

Method of administration
Voriconazole requires reconstitution and dilution (see section 6.6) prior to administration as an intravenous infusion. Not for bolus injection.

4.3. Contraindications

Hypersensitivity to the active substance or to any of the excipients listed in the formulation.
Coadministration with CYP3A4 substrates, terfenadine, astemizole, cisapride or ivabradine since increased plasma concentrations of these medicinal products can lead to QTc prolongation and rare occurrences of torsades de pointes.
Coadministration with rifampicin, carbamazepine and phenobarbital since these medicinal products are likely to decrease plasma voriconazole concentrations significantly.
Coadministration of standard doses of voriconazole with efavirenz doses of 400 mg once daily or higher is contraindicated, because efavirenz significantly decreases plasma voriconazole concentrations in healthy subjects at these doses. Voriconazole also significantly increases efavirenz plasma concentrations.
Coadministration with high-dose ritonavir (400 mg and above twice daily) because ritonavir significantly decreases plasma voriconazole concentrations in healthy subjects at this dose.
Coadministration with ergot alkaloids (ergotamine, dihydroergotamine), which are CYP3A4 substrates, since increased plasma concentrations of these medicinal products can lead to ergotism.
Coadministration with sirolimus since voriconazole is likely to increase plasma concentrations of sirolimus significantly.
Coadministration with St. John's Wort.
Coadministration with venlafaxine at initiation and during venlafaxine dose titration phase since voriconazole is likely to significantly increase plasma concentrations of venlafaxine and increase risk of tumour lysis syndrome.

4.4. Special Warnings and Precautions for Use

Hypersensitivity
Caution should be used in prescribing VFEND to patients with hypersensitivity to other azoles.

Duration of IV treatment
The duration of treatment with the intravenous formulation should be no longer than 6 months (see section 5.3).

Cardiovascular
Voriconazole has been associated with QTc interval prolongation. There have been rare cases of torsades de pointes in patients taking voriconazole who had risk factors, such as history of cardiotoxic chemotherapy, cardiomyopathy, hypokalaemia and concomitant medical products that may have been contributory. Voriconazole should be administered with caution to patients with potentially proarrhythmic conditions, such as:

- Congenital or acquired QTc prolongation.
- Cardiomyopathy, in particular when heart failure is present.
- Sinus bradycardia.
- Existing symptomatic arrhythmias.
- Concomitant medical product that is known to prolong QTc interval. Electrolyte disturbances such as hypokalaemia, hypomagnesaemia and hypocalcaemia should be monitored and corrected, if necessary, prior to initiation and during voriconazole therapy (see section 4.2). A study has been conducted in healthy volunteers which examined the effect on QTc interval of single doses of voriconazole up to 4 times the usual daily dose. No subject experienced an interval exceeding the potentially clinically-relevant threshold of 500 msec.

Infusion-related reactions
Infusion-related reactions, predominantly flushing and nausea, have been observed during administration of the intravenous formulation of voriconazole. Depending on the severity of symptoms, consideration should be given to stopping treatment.

Hepatic toxicity
In clinical trials, there have been cases of serious hepatic reactions during treatment with voriconazole (including clinical hepatitis, cholestasis and fulminant hepatic failure, including fatalities). Instances of hepatic reactions were noted to occur primarily in patients with serious underlying medical conditions (predominantly haematological malignancy). Transient hepatic reactions, including hepatitis and jaundice, have occurred among patients with no other identifiable risk factors. Liver dysfunction was generally reversible on discontinuation of therapy.

Monitoring of hepatic function
Patients receiving Voriconazole must be carefully monitored for hepatic toxicity. Clinical management should include laboratory evaluation of hepatic function (specifically AST and ALT) at the initiation of treatment with Voriconazole and at least weekly for the first month of treatment. Treatment duration should be as short as possible, however the risk/benefit assessment for treatment extension should be performed on a case-by-case basis (see section 4.2), monitoring frequency can be reduced to monthly if there are no changes in the Liver Function Tests.
If the liver function tests become markedly elevated, Voriconazole should be discontinued, unless the medical judgment of the risk-benefit of the treatment for the patient justifies continued use.
Monitoring of hepatic function should be carried out in both children and adults.

Serious dermatological adverse reactions
- Photosensitivity

In addition VFEND has been associated with photosensitivity including reactions such as epithelioma, lentigo, actinic keratosis and pseudoporphyria. It is recommended that all patients, including children, should have exposure to direct sunlight during VFEND treatment and use measures such as protective clothing and sunscreen with high sun protection factor (SPF).

- Squamous cell carcinoma of the skin

Squamous cell carcinoma of the skin has been reported in patients, some of whom have reported prior phototoxic reactions. If phototoxic reactions occur, multidisciplinary advice should be sought. If VFEND discontinuation and use of alternative antifungal agents should be considered and the patient should be referred to a dermatologist. If VFEND is continued, however, dermatologic evaluation should be performed on a systematic and regular basis, to allow early detection and management of squamous cell carcinoma. VFEND should be discontinued if premalignant skin lesions or squamous cell carcinoma are identified (see below the section under Long-term treatment).

- Severe cutaneous adverse reactions

Severe cutaneous adverse reactions (SCARs) including Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN), and drug reaction with eosinophilia and systemic symptoms (DRESS), which can be life-threatening or fatal, have been reported with the use of voriconazole. If a patient develops a rash he should be monitored closely and VFEND discontinued if lesions progress.

Adrenal events
Adrenal insufficiency has been reported in patients receiving other azoles (e.g., ketoconazole).
Reversible cases of adrenal insufficiency have been reported in patients receiving voriconazole.
Patients on long-term treatment with voriconazole (including corticosteroids including inhaled corticosteroids e.g., budesonide and intranasal corticosteroids) should be carefully monitored for adrenal insufficiency. Long-term treatment with voriconazole in the presence of corticosteroids should be discontinued.

Long-term treatment
Long-term exposure (treatment or prophylaxis) greater than 180 days (6 months) requires careful assessment of the benefit-risk balance and physicians should therefore consider the need to limit the exposure to Voriconazole.

Squamous cell carcinoma of the skin (SCC) has been reported in relation with long-term VFEND treatment.
Non-infectious periostitis with elevated fluoride and alkaline phosphatase levels has been reported in long-term patients. If a patient develops skeletal pain and radiologic findings compatible with periostitis VFEND discontinuation should be considered after multidisciplinary advice.

Visual adverse reactions
There have been reports of prolonged visual adverse reactions, including blurred vision, optic neuritis and papilloedema.

Renal adverse reactions
Acute renal failure has been observed in severely ill patients undergoing treatment with VFEND. Patients being treated with voriconazole are likely to be treated concomitantly with nephrotoxic medicinal products and have concurrent conditions that may result in decreased renal function (see section 4.8).

Monitoring of renal function
Patients should be monitored for the development of abnormal renal function. This should include laboratory evaluation, particularly serum creatinine.

Monitoring of pancreatic function
Patients, especially children, with risk factors for acute pancreatitis (e.g., recent chemotherapy, haematopoietic stem cell transplantation [HSCT]), should be monitored closely during VFEND treatment. Monitoring of serum amylase or lipase may be considered in this clinical situation.

Paediatric population
Safety and effectiveness in paediatric subjects below the age of two years has not been established (see sections 4.8 and 5.1). Voriconazole is indicated for paediatric patients aged two years or older. A higher frequency of liver enzyme elevations was observed in the paediatric population (see section 4.8). Hepatic function should be monitored in both children and adults. Oral bioavailability may be limited in paediatric patients aged 2 to <12 years with malabsorption and very low body weight for age. In that case, intravenous voriconazole administration is recommended.

- Serious dermatological adverse reactions (including SCC)

The frequency of photosensitivity reactions is higher in the paediatric population. As an evolution towards SCC has been reported, stringent measures for the photoprotection are warranted in this population of patients. In children experiencing photosensitivity reactions such as lentigines or ephelides, avoidance and dermatologic follow-up are recommended even after treatment discontinuation.
Prophylaxis in case of treatment-related adverse events (including photostability and SCC, severe or irreversible photosensitivity, or secondary invasive cancer or pre-existing photosensitivity) discontinuation of voriconazole and use of alternative antifungal agents must be considered. Phenytoin (CYP2C9 substrate and potent CYP450 inducer).

Careful monitoring of phenytoin levels is recommended when phenytoin is coadministered with voriconazole. Concomitant use of voriconazole and phenytoin should be avoided unless the benefit outweighs the risk.

Efavirenz (CYP450 inducer; CYP3A4 inhibitor and substrate)
When voriconazole is coadministered with efavirenz doses of voriconazole should be increased to 400 mg every 12 hours and the dose of efavirenz should be decreased to 300 mg every 24 hours (see sections 4.2, 4.3 and 4.5).

Rifabutin (Potent CYP450 inducer)
Careful monitoring of full blood counts and adverse reactions to rifabutin (e.g., uveitis) is recommended when rifabutin is coadministered with voriconazole. Concomitant use of voriconazole and rifabutin should be avoided unless the benefit outweighs the risk (see section 4.5).

Ritonavir (potent CYP450 inducer; CYP3A4 inhibitor and substrate)
Coadministration of voriconazole and low-dose ritonavir (100 mg twice daily) should be avoided unless an assessment of the benefit/risk to the patient justifies the use of voriconazole (see sections 4.3 and 4.5).
Everolimus (CYP3A4 substrate, P-gp substrate) Coadministration of voriconazole with everolimus is not recommended because voriconazole is expected to significantly increase everolimus concentrations. Currently there are insufficient data to allow dosing recommendations in this situation (see section 4.5).

Naloxegol (CYP3A4 substrate)
Coadministration of voriconazole and naloxegol is not recommended because voriconazole is expected to significantly increase naloxegol concentrations. Currently there are insufficient data to allow dosing recommendations in this situation (see section 4.5).

Methadone (CYP3A4 substrate)
Frequent monitoring for adverse reactions and toxicity related to methadone, including QTc prolongation, is recommended when coadministered with voriconazole since methadone levels increased following coadministration of voriconazole. Dose reduction of methadone may be needed (see section 4.5).

Short-acting opiates (CYP3A4 substrate)
Reduction in the dose of alfentanil, fentanyl and other short-acting opiates similar in structure to alfentanil and metabolised by CYP3A4 (e.g., sufentanil) should be considered when coadministered with voriconazole (see section 4.5). As the half-life of alfentanil is prolonged in a 4-fold manner when alfentanil is coadministered with voriconazole, and in an independent published study voriconazole and a single dose of alfentanil resulted in an increase in the mean AUC0–∞ of fentanyl, frequent monitoring for opiate-associated adverse reactions (including a longer respiratory monitoring period) may be necessary.

Long-acting opiates (CYP3A4 substrate)
Reduction in the dose of oxycodone and other long-acting opiates metabolized by CYP3A4 (e.g., hydrocodone) should be considered when coadministered with voriconazole. Frequent monitoring for opiate-associated adverse reactions may be necessary (see section 4.5).

Fluconazole (CYP2C9, CYP2C19 and CYP3A4 inhibitor)
Coadministration of oral voriconazole and oral fluconazole resulted in a significant increase in Cmax and AUC of voriconazole in healthy subjects. The reduced dose and/or frequency of voriconazole administration should be considered if oral azole medicinal products are used concomitantly. Monitoring for voriconazole -associated adverse reactions is recommended if voriconazole is used sequentially after fluconazole (see section 4.5).

Sodium content
This medicinal product contains 216-228 mg of sodium equivalent to 10.8-11.4% of the WHO recommended maximum daily intake of 2g sodium for an adult.

4.5. Drug Interactions

Voriconazole is metabolised by, and inhibits the activity of, cytochrome P450 isoenzymes, CYP2C19, CYP2C9, and CYP3A4. Inhibitors or inducers of these isoenzymes may increase or decrease the plasma concentrations of voriconazole, respectively. There is potential for voriconazole to increase the plasma concentrations of substances metabolised by these CYP450 isoenzymes.
Unless otherwise specified, drug interaction studies have been performed in healthy adult male subjects using multiple dosing to steady state with oral voriconazole at 200 mg twice daily (BID). These results are relevant to other populations and routes of administration.
Voriconazole should be administered with concomitant medications in patients with consideration of the potential for altering QTc interval. When there is also a potential for voriconazole to increase the plasma concentrations of substances metabolised by CYP3A4 isoenzymes (certain antihistamines, quinidine, cisapride, pimozide), co-administration is contraindicated (see below and section 4.3).

Interaction table
Interactions between voriconazole and other medicinal products are listed in the table below (once daily as "QD", twice daily as "BID", three times daily as "TID" and not determined as "ND"). The direction of the arrow for each pharmacokinetic parameter is based on the 90% confidence interval of the geometric mean ratio being within (↔), below (↓) or above (↑) the 80-125% range. The abbreviations, if included, are: AUC, AUC and AUC$_{\tau}$ represent area under the curve over a dosing interval, from time zero to the time with detectable measurement and from time zero to infinity, respectively.
The interactions in the table are presented in the following order: contraindications, those requiring dose adjustment and careful clinical and/or biological monitoring, and finally those for which no clinical information is available or that are of limited clinical significance.

Medicinal product [Mechanism of interaction]	**Interaction Geometric mean changes (%)**	**Recommendations concerning co-administration**
Anticoagulants Warfarin (30 mg single dose, co- administered with 300 mg BID voriconazole) (*CYP2C9 substrate*)	Maximum increase in prothrombin time was approximately 2-fold.	Close monitoring of prothrombin time or other suitable anticoagulation tests is recommended, and the dose of anticoagulants should be adjusted accordingly.
Other oral coumarins (e.g., phenprocoumon, acenocoumarol) (*CYP2C9 and CYP3A4 substrates*)	Although not studied, voriconazole may increase the plasma concentrations of coumarins that may cause an increase in prothrombin time.	
Benzodiazepines (e.g., midazolam, triazolam, alprazolam) (*CYP3A4 substrates*)	Although not studied clinically, voriconazole is likely to increase the plasma concentrations of benzodiazepines that are metabolised by CYP-P3A4 and lead to a prolonged sedative effect.	Dose reduction of benzodiazepines should be considered.
Immunosuppressants (*CYP3A4 substrates*) Sirolimus (2 mg single dose)	In an independent published study, Sirolimus C_{max} ↑ 6.6-fold Sirolimus AUC_{τ} ↑ 11-fold Cyclosporin C_{max} ↑ 13% Cyclosporin AUC_{τ} ↑ 70%	Co-administration of voriconazole and sirolimus is **contraindicated** (see section 4.3). When initiating voriconazole in patients already on cyclosporin it is recommended that the cyclosporin dose be halved and cyclosporin level carefully monitored. Increased cyclosporin levels have been associated with nephrotoxicity. When voriconazole is discontinued, cyclosporin levels must be carefully monitored and the dose increased as necessary. When initiating voriconazole in patients already on tacrolimus, it is recommended that the tacrolimus dose be reduced to a third of the original dose and tacrolimus level carefully monitored. Increased tacrolimus levels have been associated with nephrotoxicity. When voriconazole is discontinued, tacrolimus levels must be carefully monitored and the dose increased as necessary.
Cyclosporin (in stable renal transplant recipients receiving chronic cyclosporin therapy)		
Tacrolimus (5 mg/kg single dose)	Tacrolimus C_{max} ↑ 117% Tacrolimus AUC_{τ} ↑ 221%	
Long-Acting Opiates (*CYP3A4 substrates*) Oxycodone (10 mg single dose)	In an independent published study, Oxycodone C_{max} ↑ 1.7-fold Oxycodone AUC_{τ} ↑ 3.6-fold	Dose reduction in oxycodone and other long-acting opiates metabolized by CYP3A4 (e.g., hydrocodone) should be considered. Frequent monitoring for opiate-associated adverse reactions may be necessary (see section 4.4).
Methadone (32-100 mg) (QD) (*CYP3A4 substrate*)	R-methadone (active) C_{max} ↑ 31% R-methadone (active) AUC_{τ} ↑ 47% S-methadone C_{max} ↑ 65% S-methadone AUC_{τ} ↑ 103%	Frequent monitoring for adverse reactions and toxicity related to methadone, including QTc prolongation, is recommended. Dose reduction of methadone may be needed.
Non-Steroidal Anti-inflammatory Drugs (NSAIDs) (*CYP2C9 substrates*) Ibuprofen (400 mg single dose) Diclofenac (50 mg single dose)	S-Ibuprofen C_{max} ↑ 20% S-Ibuprofen AUC_{τ} ↑ 100% Diclofenac C_{max} ↑ 114% Diclofenac AUC_{τ} ↑ 78%	Frequent monitoring for adverse reactions and toxicity related to NSAIDs is recommended. Dose reduction of NSAIDs may be needed.
Omeprazole (40 mg QD)* (*CYP2C19 inhibitor; CYP2C19 and CYP3A4 substrate*)	Omeprazole C_{max} ↑ 116% Omeprazole AUC_{τ} ↑ 280% Omeprazole C_{max} ↑ 10% Omeprazole AUC_{τ} ↑ 15% Other proton pump inhibitors that are CYP2C19 substrates may also be inhibited by voriconazole and may result in increased plasma concentrations of these medicinal products.	No dose adjustment of voriconazole is recommended. When initiating voriconazole in patients already receiving omeprazole at 40 mg or above, it is recommended that the omeprazole dose be halved.
Oral Contraceptives* (*CYP3A4 substrate; CYP2C19 inhibitor*) Norethisterone/ethinylestradiol (1 mg/0.035 mg QD)	Ethinylestradiol C_{max} ↑ 36% Ethinylestradiol AUC_{τ} ↑ 61% Norethisterone C_{max} ↑ 15% Norethisterone AUC_{τ} ↑ 53% Voriconazole C_{max} ↓ 18% Voriconazole AUC_{τ} ↑ 46%	Monitoring for adverse reactions related to oral contraceptives, in addition to those for voriconazole, is recommended.
Short-acting Opiates (*CYP3A4 substrates*) Alfentanil (20 µg/kg single dose, with concomitant naloxone) Fentanyl (5 µg/kg single dose)	In an independent published study, Alfentanil AUC_{τ} ↑ 6-fold In an independent published study, Fentanyl AUC_{τ} ↑ 1.34-fold	Dose reduction of alfentanil, fentanyl and other short-acting opiates similar in structure to alfentanil and metabolised by CYP3A4 (e.g., sufentanil) should be considered. Extended and frequent monitoring for respiratory depression and other opiate-associated adverse reactions is recommended.
Statins (e.g., lovastatin) (*CYP3A4 substrates*)	Although not studied clinically, voriconazole is likely to increase the plasma concentrations of statins that are metabolised by CYP3A4 and lead to a prolonged effect.	Dose reduction of statins should be considered.
Sulfonylureas (e.g., tolbutamide, glipizide, glyburide) (*CYP2C9 substrates*)	Although not studied, voriconazole is likely to increase the plasma concentrations of sulfonylureas and cause hypoglycaemia.	Careful monitoring of blood glucose is recommended and dose reduction of sulfonylureas should be considered.
Vinca Alkaloids (e.g., vincristine and vinblastine) (*CYP3A4 substrates*)	Although not studied, voriconazole is likely to increase the plasma concentrations of vinca alkaloids and lead to neurotoxicity.	Dose reduction of vinca alkaloids should be considered.
Other HIV Protease Inhibitors (e.g., saquinavir, amprenavir and nelfinavir)* (*CYP3A4 substrates and inhibitors*)	Not studied clinically. *In vitro* studies show that voriconazole may inhibit the metabolism of HIV protease inhibitors and, reciprocally, the latter may also be inhibited by HIV protease inhibitors.	Careful monitoring for any occurrence of drug toxicity and/or lack of efficacy, and dose adjustment may be needed.
Other Non-Nucleoside Reverse Transcriptase Inhibitors (NNRTIs) (e.g., delavirdine, nevirapine)* (*CYP3A4 substrates, inhibitors or CYP450 inducers*)	Not studied clinically. *In vitro* studies show that the metabolism of voriconazole may be inhibited by NNRTIs and voriconazole may inhibit the metabolism of NNRTIs. The findings of the effect of efavirenz on voriconazole suggest that the metabolism of voriconazole may be induced by an NNRTI.	Careful monitoring for any occurrence of drug toxicity and/or lack of efficacy, and dose adjustment may be needed.
Cimetidine (400 mg BID) (*non-specific CYP450 inhibitor and increases gastric pH*)	Voriconazole C_{max} ↑ 18% Voriconazole AUC_{τ} ↑ 23%	No dose adjustment
Digoxin (0.25 mg QD) (*P-gp substrate*)	Digoxin C_{max} ↔ Digoxin AUC_{τ} ↔	No dose adjustment
Indinavir (800 mg TID) (*CYP3A4 inhibitor and substrate*)	Indinavir C_{max} ↔ Indinavir AUC_{τ} ↔ Voriconazole C_{max} ↔ Voriconazole AUC_{τ} ↔	No dose adjustment
Macrolide antibiotics Erythromycin (1 g BID) (*CYP3A4 substrate*) Azithromycin (500 mg QD)	Voriconazole C_{max} and AUC_{τ} ↔ Voriconazole C_{max} and AUC_{τ} ↔ The effect of voriconazole on either erythromycin or azithromycin is unknown.	No dose adjustment
Mycophenolic acid (1 g single dose) (*UDP-glucuronyl transferase substrate*)	Mycophenolic acid C_{max} ↔ Mycophenolic acid AUC_{τ} ↔	No dose adjustment
Prednisolone (60 mg single dose) (*CYP3A4 substrate*)	Prednisolone C_{max} ↑ 11% Prednisolone AUC_{τ} ↑ 34%	No dose adjustment
Ranitidine (150 mg BID) (*increases gastric pH*)	Voriconazole C_{max} and AUC_{τ} ↔	No dose adjustment

4.6. Use in Special Populations (such as pregnant women, lactating women, paediatric patients, geriatric patients etc.)

Pregnancy
There are no adequate data on the use of Voriconazole in pregnant women.
Studies in animals have shown reproductive toxicity (see section 5.3). The potential risk for humans is unknown.
Voriconazole must not be used during pregnancy unless the benefit to the mother clearly outweighs the potential risk to the foetus.

Women of Child-bearing potential
Women of child-bearing potential must always use effective contraception during treatment.

Breast-feeding
The excretion of voriconazole into breast milk has not been investigated. Breast-feeding must be stopped on initiation of treatment with Voriconazole.

Fertility
In an animal study, no impairment of fertility was demonstrated in male and female rats (see section 5.3).

Elderly
No dose adjustment is necessary for elderly patients (see section 5.2).

Renal impairment
Dosage adjustments to severe renal dysfunction (creatinine clearance < 50 ml/min), accumulation of the intravenous vehicle, SBECD, occurs. Oral voriconazole should be administered to these patients, unless an assessment of the risk benefit to the patient justifies the use of intravenous voriconazole. Serum creatinine levels should be monitored and if there is evidence of deterioration in renal function, consideration should be given to oral voriconazole therapy (see section 5.2).
Voriconazole is haemodialysed with a clearance of 121 ml/min. A 4 hour haemodialysis session does not remove a sufficient amount of voriconazole to warrant dose adjustment.
The intravenous vehicle, SBECD, is haemodialysed with a clearance of 55 ml/min.

Hepatic impairment
It is recommended that the standard loading dose regimen be used but that the maintenance dose be halved in patients with mild to moderate hepatic cirrhosis (Child-Pugh A and B) receiving voriconazole (see section 5.2).
No dose adjustment has been studied in patients with severe hepatic cirrhosis and if necessary close monitoring for drug toxicity (see section 4.8).
There is limited data on the safety of voriconazole in patients with abnormal Liver Function Tests (aspartate transaminase [AST], alanine transaminase [ALT], alkaline phosphatase [ALP]) or bilirubin >5 times the upper limit of normal.
Voriconazole has been associated with elevations in liver function tests and clinical signs of hepatic cause, such as jaundice, and must only be used in patients with severe hepatic impairment if the benefit outweighs the potential risk. Patients with severe hepatic impairment must be carefully monitored for drug toxicity (see section 4.8).

Paediatric population
The safety and efficacy of voriconazole in children below 2 years has not been established.

4.7. Effects on Ability to Drive and Use Machines

Voriconazole has moderate influence on the ability to drive and use machines. It may cause transient and reversible changes to vision, including blurring, photophobia and/or increased perception and/or photophobia. Patients must avoid potentially hazardous tasks, such as driving or operating machinery, while experiencing these symptoms.

4.8. Undesirable Effects

The adverse reactions of Voriconazole are presented in below table. The adverse reactions are listed by system organ class (SOC) and frequency. Frequency categories are defined as: Very common (≥1/10); Common (≥1/100 to <1/10); Uncommon (≥1/1,000 to <1/100); Rare (≥1/10,000 to <1/1,000); Very rare (<1/10,000); Not known (cannot be estimated from the available data).
Within each frequency grouping, undesirable effects are presented in order of decreasing seriousness.

Undesirable effects reported in subjects receiving voriconazole:

System Organ Class	Very common	Common	Uncommon	Rare	Frequency not known
Infections and infestations		sinusitis	pseudomembranous colitis		
Neoplasms benign, malignant and unspecified (including cysts and polyps)					squamous cell carcinoma*
Blood and lymphatic system disorders		agranulocytosis†, pancytopenia, thrombocytopenia†, leukopenia, anaemia	bone marrow failure, lymphadenopathy, eosinophilia	disseminated intravascular coagulation	
Immune system disorders			hypersensitivity	anaphylactoid reaction	
Endocrine disorders			adrenal insufficiency, hypothyroidism	hyperthyroidism	
Metabolism and nutrition disorders	oedema peripheral	hypoglycaemia, hypokalaemia, hyponatraemia			
Psychiatric disorders		depression, hallucination, anxiety, confusion, agitation, confusional state			
Nervous system disorders	headache		brain oedema, encephalopathy*, dysgeusia, hepatic encephalopathy, dizziness, hypoaesthesia, somnolence, tremor	hepatic encephalopathy, Guillain-Barre syndrome, nystagmus	
Eye disorders	visual impairment*	retinal haemorrhage	optic neuritis, papilloedema	optic atrophy, corneal opacity	
Ear and labyrinth disorders			hypoacusis, vertigo, tinnitus		
Cardiac disorders			ventricular fibrillation, ventricular extrasystoles, ventricular tachycardia, electrocardiogram QT prolonged, supraventricular tachycardia	torsades de pointes, atrioventricular block complete, bundle branch block, nodal rhythm	
Vascular disorders		hypotension, phlebitis			
Respiratory, thoracic and mediastinal disorders	respiratory distress*	respiratory distress syndrome, pulmonary oedema			
Gastrointestinal disorders	diarrhoea, vomiting, abdominal pain, nausea	cheilitis, dyspepsia, constipation, glossitis	pancreatitis, pancreatitis*, stomatitis, duodenitis, gastroenteritis, gingivitis		
Hepatobiliary disorders	liver function test abnormal	jaundice, jaundice cholestatic, hepatitis†	hepatic failure, cholangitis, cholestasis, cholelithiasis		
Skin and subcutaneous tissue disorders	rash	dermatitis exfoliative, alopecia, maculo-papular rash, pruritus, erythema	Stevens-Johnson syndrome, photosensitivity, psoriasis, dermatitis allergic, exfoliative skin disorder, rash pustular, rash macular, eczema	toxic epidermal necrolysis, drug reaction with eosinophilia and systemic symptoms (DRESS)†, angioedema, actinic keratosis*, pseudoporphyria, erythema multiforme, psoriasis, drug eruption	cutaneous lupus erythematosus*, epithelioma*, lentigo*
Musculoskeletal and connective tissue disorders		back pain	arthritis		periostitis*
Renal and urinary disorders		renal failure acute, haematuria	renal tubular necrosis, proteinuria, nephritis		
General disorders and administration site conditions	pyrexia	chest pain, face oedema*, asthenia, chills	infusion site reaction, influenza like illness		
Investigations		blood creatinine increased	blood urea increased, blood cholesterol increased		

* ADR identified post-marketing
† Includes life-threatening and neutropenia.
‡ Includes immune thrombocytopenic purpura.
§ Includes drug hypersensitivity.
¶ Includes hypoxic-ischaemic encephalopathy and metabolic encephalopathy.
# Includes abathosis and parakeratosis.
* See "Visual impairments" paragraph in section 4.8.
** See "Hepatic toxicity" paragraph post-marketing. See section 4.4.
†† See section 4.4.
‡‡ Includes dyspnoea and dyspnoea exertional.
§§ Includes drug-induced liver injury, hepatitis toxic, hepatocellular injury and hepatotoxicity.
¶¶ Includes periorbital oedema, lip oedema, and oedema mouth.

Reporting of suspected adverse reactions
Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. To report Suspected Adverse Reactions, contact MSN Laboratories Private Limited at pharmacovigilance@msnlabs.com or through company website www.msnlabs.com->Contact us->Medical Enquiry/ to report a side effect.

4.9. Overdose

In clinical trials there were 3 cases of accidental overdose. All occurred in paediatric patients, who received up to five times the recommended intravenous dose of voriconazole. A single adverse reaction of photophobia of 10 minutes duration was reported.
There is no known antidote to voriconazole.
Voriconazole is haemodialysed with a clearance of 121 ml/min. The intravenous vehicle, SBECD, is haemodialysed with a clearance of 55 ml/min. In an overdose, haemodialysis may assist in the removal of voriconazole and SBECD from the body.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Antimycotics for systemic use, triazole derivatives, ATC code: J02 AC03

Mode of action
Voriconazole is a triazole antifungal agent. The primary mode of action of voriconazole is the inhibition of fungal cytochrome P450-mediated 14 alpha-lanosterol demethylation, an essential step in fungal ergosterol biosynthesis. The accumulation of 14 alpha-methyl sterols correlates with the subsequent loss of ergosterol in the fungal cell membrane and is thought to be responsible for the antifungal activity of voriconazole. Voriconazole has been shown to be more selective for fungal cytochrome P-450 enzymes than for various mammalian cytochrome P-450 enzyme systems.

Pharmacokinetic-pharmacodynamic relationship
In 10 therapeutic studies, the median for the average and maximum plasma concentrations in individual subjects across the studies was 2425 ng/ml and 5 (inter-quartile range: 1193 to 4380 ng/ml) and