

**WARNING:** To be sold by retail on the prescription of Gastroenterologist/Hepatologist only.

R

## Trentine Hydrochloride Capsules USP 250 mg

### Cuchel एम एस एस व्हार्फेल

#### Label Claim

Each hard gelatin capsule contains

Trentine Hydrochloride USP 250 mg

#### Excipients Q.S & Approved colors used in capsule shell

Dosage form/strength : capsule: 250 mg

**Chemical Name:** Trentine hydrochloride is *N,N*-bis (2-aminoethyl)-1,2-ethanediamine dihydrochloride. It is a white to pale yellow crystalline hygroscopic powder. It is freely soluble in water, soluble in methanol, slightly soluble in ethanol, and insoluble in chloroform and ether.

**Structural formula is:** NH<sub>2</sub>(CH<sub>2</sub>)<sub>2</sub>NH(CH<sub>2</sub>)<sub>2</sub>NH<sub>2</sub>·2HCl

**Empirical formula :** C<sub>6</sub>H<sub>18</sub>N<sub>4</sub>·2HCl

**Molecular weight :** 219.2

**Route of administration :** oral

#### INDICATIONS AND USAGE :

Trentine Hydrochloride Capsules are indicated in the treatment of patients with Wilson's disease who are intolerant of penicillamine. Clinical experience with Trentine Hydrochloride Capsules is limited and alternate dosing regimens have not been well characterized; all endpoints in determining an individual patient's dose have not been well defined. Trentine Hydrochloride Capsules and penicillamine cannot be considered interchangeable. Trentine Hydrochloride Capsules should be used when continued treatment with penicillamine is no longer possible because of intolerance or life endangering side effects.

Unlike penicillamine, Trentine Hydrochloride Capsules are not recommended in cystinuria or rheumatoid arthritis. The absence of a sulphydryl moiety renders it incapable of binding cystine and, therefore, it is of no use in cystinuria. In 15 patients with rheumatoid arthritis, Trentine Hydrochloride Capsules were reported not to be effective in improving any clinical or biochemical parameter after 12 weeks of treatment.

Trentine Hydrochloride Capsules are not indicated for treatment of biliary cirrhosis.

#### DOSAGE AND ADMINISTRATION

Systemic evaluation of dose and/or interval between dose has not been done. However, on limited clinical experience, the recommended initial dose of Trentine Hydrochloride Capsules is 500-750 mg/day for pediatric patients and 750-1250 mg/day for adults given in divided doses two, three or four times daily. This may be increased to a maximum of 2000 mg/day for adults or 1500 mg/day for pediatric patients age 12 or under.

The daily dose of Trentine Hydrochloride Capsules should be increased only when the clinical response is not adequate or the concentration of free serum copper is persistently above 20 mcg/dL. Optimal long-term maintenance dosage should be determined at 6-12 month intervals.

It is important that Trentine Hydrochloride Capsules be given on an empty stomach, at least one hour before meals or two hours after meals and at least one hour apart from any other drug, food, or milk. The capsules should be swallowed whole with water and should not be opened or chewed.

#### CONTRAINDICATIONS

Hypersensitivity to this product.

#### WARNINGS

Patient experience with trentine hydrochloride is limited (see **CLINICAL PHARMACOLOGY**). Patients receiving Trentine Hydrochloride Capsules should remain under regular medical supervision throughout the period of drug administration. Patients (especially women) should be closely monitored for evidence of iron deficiency anemia.

#### PRECAUTIONS

##### General

There are no reports of hypersensitivity in patients who have been administered trentine hydrochloride for Wilson's disease. However, there have been reports of asthma, bronchitis and dermatitis occurring after prolonged environmental exposure in workers who use trentine hydrochloride as a hardener of epoxy resins. Patients should be observed closely for signs of possible hypersensitivity.

##### Information for Patients

Patients should be directed to take Trentine Hydrochloride Capsules on an empty stomach, at least one hour before meals or two hours after meals and at least one hour apart from any other drug, food, or milk. The capsules should be swallowed whole with water and should not be opened or chewed. Because of the potential for contact dermatitis, any site of exposure to the capsule contents should be washed with water promptly. For the first month of treatment, the patient should have his temperature taken nightly, and he should be asked to report any symptom such as fever or skin eruption.

##### Laboratory Tests

The most reliable index for monitoring treatment is the determination of free copper in the serum, which equals the difference between quantitatively determined total copper and ceruloplasmin-copper. Adequately treated patients will usually have less than 10 mcg free copper/dL of serum.

Therapy may be monitored with a 24-hour urinary copper analysis periodically (i.e., every 6-12 months). Urine must be collected in copper-free glassware. Since a low copper diet should keep copper absorption down to less than one milligram a day, the patient probably will be in the desired state of negative copper balance if 0.5 to 1.0 milligram of copper is present in a 24-hour collection of urine.

##### Drug Interactions

In general, mineral supplements should not be given since they may block the absorption of Trentine Hydrochloride Capsules. However, iron deficiency may develop, especially in children and menstruating or pregnant women, or as a result of the low copper diet recommended for Wilson's disease. If necessary, iron may be given in short courses, but since iron and Trentine Hydrochloride Capsules each inhibit absorption of the other, two hours should elapse between administration of Trentine Hydrochloride Capsules and iron.

It is important that Trentine Hydrochloride Capsules be taken on an empty stomach, at least one hour before meals or two hours after meals and at least one hour apart from any other drug, food, or milk. This permits maximum absorption and reduces the likelihood of inactivation of the drug by metal binding in the gastrointestinal tract.

##### Carcinogenesis, Mutagenesis, Impairment of Fertility

Data on carcinogenesis, mutagenesis, and impairment of fertility are not available.

##### Pregnancy

Trentine hydrochloride was teratogenic in rats at doses similar to the human dose. The frequencies of both resorptions and fetal abnormalities, including hemorrhage and edema, increased while fetal copper levels decreased when trentine hydrochloride was given in the maternal diets of rats. There are no adequate and well-controlled studies in pregnant women. Trentine Hydrochloride Capsules should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

##### Nursing Mothers

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when Trentine Hydrochloride Capsules are administered to a nursing mother.

##### Pediatric Use

Controlled studies of the safety and effectiveness of Trentine Hydrochloride Capsules in pediatric patients have not been conducted. It has been used clinically in pediatric patients as young as 6 years with no reported adverse experiences.

##### Geriatric Use

Clinical studies of Trentine Hydrochloride Capsules did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience is insufficient

to determine differences in responses between the elderly and younger patients. In general, dose selection should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

#### ADVERSE REACTIONS

Clinical experience with Trentine Hydrochloride Capsules has been limited. The following adverse reactions have been reported in a clinical study in patients with Wilson's disease who were on therapy with trentine hydrochloride: iron deficiency, systemic lupus erythematosus. In addition, the following adverse reactions have been reported in marketed use: dystonia, muscular spasm, myasthenia gravis. Trentine Hydrochloride Capsules are not indicated for treatment of biliary cirrhosis, but in one study of 4 patients treated with trentine hydrochloride for primary biliary cirrhosis, the following adverse reactions were reported: heartburn; epigastric pain and tenderness; thickening, fissuring and flaking of the skin; hypochromic microcytic anemia; acute gastritis; aphthous ulcers; abdominal pain; melena; anorexia; malaise; cramps; muscle pain; weakness; rhabdomyolysis. A causal relationship of these reactions to drug therapy could not be rejected or established.

#### OVERDOSE

There is a report of an adult woman who ingested 30 grams of trentine hydrochloride without apparent ill effects. No other data on overdosage are available.

#### CLINICAL PHARMACOLOGY

##### Introduction

Wilson's disease (hepatolenticular degeneration) is an autosomal inherited metabolic defect resulting in an inability to maintain a near-zero balance of copper. Excess copper accumulates possibly because the liver lacks the mechanism to excrete free copper into the bile. Hepatocytes store excess copper but when their capacity is exceeded copper is released into the blood and is taken up into extrahepatic sites. This condition is treated with a low copper diet and the use of chelating agents that bind copper to facilitate its excretion from the body.

##### Clinical Summary

Forty-one patients (18 male and 23 female) between the ages of 6 and 54 with a diagnosis of Wilson's disease and who were intolerant of d-penicillamine were treated in two separate studies with trentine hydrochloride. The dosage varied from 450 to 2400 mg per day. The average dosage required to achieve an optimal clinical response varied between 1000 mg and 2000 mg per day. The mean duration of trentine hydrochloride therapy was 48.7 months (range 2-164 months). Thirty-four of the 41 patients improved, 4 had no change in clinical global response, 2 were lost to follow-up and one showed deterioration in clinical condition. One of the patients who improved while on therapy with trentine hydrochloride experienced a recurrence of the symptoms of systemic lupus erythematosus which had appeared originally during therapy with penicillamine. Therapy with trentine hydrochloride was discontinued. No other adverse reactions, except iron deficiency, were noted among any of these 41 patients.

One investigator treated 13 patients with trentine hydrochloride following their development of intolerance to d-penicillamine. Retrospectively, he compared these patients to an additional group of 12 patients with Wilson's disease who were both tolerant of and controlled with d-penicillamine therapy, but who failed to continue any copper chelation therapy. The mean age at onset of disease of the latter group was 12 years as compared to 21 years for the former group. The trentine hydrochloride group received d-penicillamine for an average of 4 years as compared to an average of 10 years for the non-treated group.

Various laboratory parameters showed changes in favor of the patients treated with trentine hydrochloride. Free and total serum copper, SGOT, and serum bilirubin all showed mean increases over baseline in the untreated group which were significantly larger than with the patients treated with trentine hydrochloride. In the 13 patients treated with trentine hydrochloride, previous symptoms and signs relating to d-penicillamine intolerance disappeared in 8 patients, improved in 4 patients, and remained unchanged in one patient. The neurological status in the trentine hydrochloride group was unchanged or improved over baseline, whereas in the untreated group, 6 patients remained unchanged and 6 worsened. Kayser-Fleischer rings improved significantly during trentine hydrochloride treatment.

The clinical outcome of the two groups also differed markedly. Of the 13 patients on therapy with trentine hydrochloride (mean duration of therapy 4.1 years; range 1 to 13 years), all were alive at the data cutoff date, and in the non-treated group (mean years with no therapy 2.7 years; range 3 months to 9 years), 9 of the 12 died of hepatic disease.

##### Chelating Properties

##### Preclinical Studies

Studies in animals have shown that trentine hydrochloride has cupriuretic activities in both normal and copper-loaded rats. In general, the effects of trentine hydrochloride on urinary copper excretion are similar to those of equimolar doses of penicillamine, although in one study they were significantly smaller.

##### Human Studies

Renal clearance studies were carried out with penicillamine and trentine hydrochloride on separate occasions in selected patients treated with penicillamine for at least one year. Six-hour excretion rates of copper were determined off treatment and after a single dose of 500 mg of penicillamine or 1.2 g of trentine hydrochloride. The mean urinary excretion rates of copper were as follows:

No. of Patients	Single Dose Treatment	Basal Excretion Rate (µg Cu ++ /6hr)	Test-dose Excretion Rate (µg Cu ++ /6hr)
1	Trentine, 1.2 g	19	234
2	Penicillamine, 500 mg	17	320

In patients not previously treated with chelating agents, a similar comparison was made:

No. of Patients	Single Dose Treatment	Basal Excretion Rate (µg Cu ++ /6hr)	Test-dose Excretion Rate (µg Cu ++ /6hr)
1	Trentine, 1.2 g	71	1326
2	Penicillamine, 500 mg	68	1074

These results demonstrate that Trentine Hydrochloride Capsules are effective as a cupriuretic agent in patients with Wilson's disease although on a molar basis it appears to be less potent or less effective than penicillamine. Evidence from a radio-labelled copper study indicates that the different cupriuretic effect between these two drugs could be due to a difference in selectivity of the drugs for different copper pools within the body.

##### Pharmacokinetics

Data on the pharmacokinetics of trentine hydrochloride are not available. Dosage adjustment recommendations are based upon clinical use of the drug.

**Storage:** Store at 2°-8°C (36°-46°F).

Keep out of reach of children.

Manufactured by:

**MSN Laboratories Private Limited,**

Formulation Division,

Unit-II, Sy.No. 1277, 1319 to 1324,

Nandigama (Village & Mandal),

Rangareddy (District),

Telangana - 509 228, India.